

can provide more data collected quickly and conveniently, which, for example, can be applied in to modelling the surface flow of a river.

REFERENCES

Bunn, S.E. and Arthington, A.H., 2002. Basic principles and ecological consequences of altered flow regimes for aquatic biodiversity, *Environmental Management*, vol. 30, no. 4, pp. 492–507.

Dobriyal, P.; Badol, R.; Tuboi, C. & Hussain, S.A., 2017. A review of methods for monitoring streamflow for sustainable water resource management, *Applied Water Science*, October 2017, Volume 7, Issue 6, pp 2617–2628 (Springer).

Helmut E., 2014. Abschluss der Renaturierung: Die Aa ist wieder ein Fluss, *Münstersche Zeitung* (6 June 2014).

Hudson, 1993. Field measurement of soil erosion and runoff, Bedford: FAO.

Jähne, B., 2005. *Digitale Bildverarbeitung*, 580p. (Springer).

Laganière, R., 2014. *OpenCV Computer Vision Application Programming Cookbook*, Second Edition, 376p. (Packt Publishing).

Matlab version 9.2.0, 2017. computer software, Massachusetts: The MathWorks Inc.

OpenCV.org, 2017. computer software, Available: opencv.org [18 Jul 2017].

Steiger, J., Tabacchi, E., Dufour, S., Corenblit, D. and Peiry, J.L., 2005. Hydrogeomorphic processes affecting riparian habitat within alluvial channel–floodplain river systems: a review for the temperate zone, *River Research and Applications*, vol. 21, no. 7, pp.719–737.

Vargas Luna, A., Crosato, A., & Uijttewaal, WSJ., 2015. Effects of vegetation on flow and sediment transport: Comparative analyses and validation of predicting models. *Earth Surface Processes and Landforms*, 40(2),157-176. <https://doi.org/10.1002/esp.3633>

Vision Wasser, 2010. Renaturierung der Münsterschen Aa nahe Nevinghoff, [Online], Available:www.visionwasser.de/kooperationen/ms/ms-aa/90.html [15 Jul 2017].